

MYCOTAXON

<http://dx.doi.org/10.5248/123.103>

Volume 123, pp. 103–106

January–March 2013

***Cordochaete* (Agaricomycetes), a new corticioid genus from India**

S.K. SANYAL, SAMITA, G.S. DHINGRA* & AVNEET P. SINGH

*Department of Botany, Punjabi University, Patiala 147 002, India**CORRESPONDENCE TO: dhingragurpaul@gmail.com

ABSTRACT – A new monotypic corticioid genus, *Cordochaete*, is described with the new species *C. cystidiata*, from Haridwar in Uttarakhand, India.

KEY WORDS – hyphal cordons, Chandighat, angiospermous wood

While conducting a fungal foray in Chandighat area in district Haridwar of Uttarakhand, India, S.K. Sanyal collected a specimen on decaying angiospermous wood in a mixed forest. The specimen was compared macroscopically and microscopically with similar genera in *Corticiaceae* (Rattan 1977, Eriksson et al. 1981, Burdsall 1985, Dhingra 1987, Nakasone & Sytsma 1993, Greslebin et al. 2004, Martínez & Nakasone 2005, Zmitrovich & Ezhov 2011) but could not be assigned to any already known, and is described here as a new genus. Morphological traits show similarities with the genera *Phlebiopsis* and *Rhizochaete*.

***Cordochaete* Sanyal, Samita, Dhingra & Avneet P. Singh, gen. nov.**

MYCOBANK 801039

Differs from *Phlebiopsis* in having hyphal cordons and is distinguished from *Rhizochaete* by the absence of a red or violet KOH reaction in both the hymenium and hyphal cordons.

TYPE SPECIES: *Cordochaete cystidiata* Sanyal et al.

ETYMOLOGY: The generic name is based on the resemblance to *Rhizochaete* and the presence of hyphal cordons.

Basidiocarp resupinate, adnate, effused; hymenial surface corneous as dry; hyphal system monomitic; generative hyphae without clamps; basal zone very thin, context conspicuous, of loosely interwoven hyphae; subhymenium compact, hyphal cordons both in context and subhymenium; cystidia thick-walled, encrusted; basidia clavate, 4-sterigmate, without basal clamp; basidiospores ellipsoidal to subcylindrical, inamyloid, acyanophilous.




PLATE 1. *Cordochaete cystidiata* (holotype). Figs 1–2: Basidiocarps. 1. Fresh. 2. Dried. Figs 3–7: Microscopic structures. 3. Basidiospores. 4. Basidia. 5. Generative hyphae. 6. Cystidia. 7. Section. Figs 8–9: Micrographs. 8. Encrusted cystidia and basidiospores. 9. Section.

REMARKS— This genus can be differentiated from *Rhizochaete* in lacking a reddish or purplish change of the basidiome and hyphal cordons in KOH solution and from *Phlebiopsis* in lacking very compact context.

Cordochaete cystidiata Sanyal, Samita, Dhingra & Avneet P. Singh, **sp. nov.**

MYCOBANK 801040

FIGS 1–9

Differs from *Phlebiopsis gigantea* by its loosely arranged context and presence of cystidia only in the hymenium and from *Rhizochaete radicata* by its compact hymenium and subhymenium.

TYPE: India, Uttarakhand: Haridwar, Chandighat, on decaying angiospermous wood, 20 August 2011, S.K. Sanyal 4745 (PUN, **holotype**).

ETYMOLOGY: The epithet refers to the presence of encrusted cystidia.

Basidiocarp resupinate, adnate, effused, $\leq 850\ \mu\text{m}$ thick in section; hymenial surface reticulate, corneous as dry, pale orange to grayish orange when fresh, orange gray to brownish gray on drying; margins thinning, fibrillose, paler concolorous to whitish, to indeterminate. Hyphal system monomitic; generative hyphae $\leq 6.5\ \mu\text{m}$ wide; generative hyphae branched, septate, without clamps; basal zone of a very few horizontal hyphae, followed by conspicuous zone of loosely interwoven hyphae interspersed with hyphal cordons; subhymenial zone mainly of compact vertical hyphae and hyphal cordons; hyphae in the cordons thick-walled, without clamps. Cystidia $85\text{--}95 \times 11\text{--}13.5\ \mu\text{m}$, subfusiform to fusiform, thick-walled, encrusted, projecting beyond the hymenial surface, negative to sulphovanillin. Basidia $28\text{--}39 \times 5\text{--}6.5\ \mu\text{m}$, clavate, 4-sterigmate, thin- to somewhat thick-walled, without basal clamp; sterigmata $\leq 5\ \mu\text{m}$ long. Basidiospores $6.5\text{--}8 \times 3\text{--}3.5\ \mu\text{m}$, ellipsoidal to subcylindrical, thin- to somewhat thick-walled, inamyloid, acyanophilous.

REMARKS— *Rhizochaete radicata* differs in having basidiome that turns reddish purple in KOH solution and subicular hyphae with brown granules that dissolve in KOH. *Phlebiopsis gigantea* differs in having cystidia distributed throughout the basidiome.

Acknowledgements

The authors thank Head, Department of Botany, Punjabi University, Patiala, for providing research facilities; Prof. Nils Hallenberg (University of Gothenburg, Gothenburg, Sweden) for expert comments and peer review; Prof. B.M. Sharma (Department of Plant Pathology, COA, CSKHPAU, Palampur, H.P., India) for peer review.

Literature cited

- Burdsall HH. 1985. A contribution to the taxonomy of the genus *Phanerochaete* (Corticaceae, Aphyllophorales). Mycologia Mem. 10. 165 p.
- Dhingra GS. 1987. The genus *Phlebiopsis* in the Eastern Himalayas. Nova Hedwigia 44: 221–227.

- Eriksson J, Hjortstam K, Ryvarden L. 1981. The *Corticiaceae* of North Europe – VI. Fungiflora, Oslo. pp. 1051–1276.
- Greslebin A, Nakasone KK, Rajchenberg M. 2004. *Rhizochaete*, new genus of phanerochaetoid fungi. *Mycologia* 96: 260–271. <http://dx.doi.org/10.2307/3762062>
- Martínez S, Nakasone KK. 2005. The genus *Phanerochaete* (*Corticiaceae*, *Basidiomycotina*) sensu lato in Uruguay. *Sydowia* 57(1): 94–101.
- Nakasone KK, Sytsma KJ. 1993. Biosystematic Studies on *Phlebia acerina*, *P. rufa*, and *P. radiata* in North America. *Mycologia* 85(6): 996–1016. <http://dx.doi.org/10.2307/3760683>
- Rattan SS. 1977. The resupinate *Aphylllophorales* of the North Western Himalayas. *Bibliotheca Mycologica* 60. 427 p.
- Zmitrovich IV, Ezhov ON. 2011. Ecology and plectology of *Phlebia tremelloidea* (*Polyporales*, *Agaricomycetes*). *Acta Mycologica* 46(1): 19–25.